

LETTER • **OPEN ACCESS**

Climate exceeded human management as the dominant control of fire at the regional scale in California's Sierra Nevada

To cite this article: Richard S Vachula *et al* 2019 *Environ. Res. Lett.* **14** 104011

View the [article online](#) for updates and enhancements.

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
28 July 2019

REVISED
19 September 2019

ACCEPTED FOR PUBLICATION
20 September 2019

PUBLISHED
15 October 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Climate exceeded human management as the dominant control of fire at the regional scale in California's Sierra Nevada

Richard S Vachula^{1,2} , James M Russell^{1,2} and Yongsong Huang^{1,2}

¹ Department of Earth, Environmental, and Planetary Sciences, Brown University, Providence, RI 02912, United States of America

² Institute at Brown for Environment and Society, Brown University, Providence, RI 02912, United States of America

E-mail: richard_vachula@brown.edu

Keywords: paleofire, fire management, climate change, paleoecology, paleoclimate

Supplementary material for this article is available [online](#)


Abstract

The societal impacts of recent, severe fires in California highlight the need to understand the long-term effectiveness of human fire management. The relative influences of local management and climate at centennial timescales are controversial and poorly understood. This is the case in California's Sierra Nevada, an actively managed area with a rich history of Native American fire use. We analyzed charcoal preserved in lake sediments from Yosemite National Park and spanning the last 1400 years to reconstruct local and regional area burned. Warm and dry climates promoted burning at both local and regional scales. However, at local scales fire management by Native Americans before 850 and between ca. 1350 and 1600 CE and, subsequently, Yosemite park managers from ca. 1900 to 1970 CE, decoupled fire extent dictated by regional climate scenarios. Climate acts as a top-down, broader scale control of fire, but human management serves a bottom-up, local control. Regional area burned peaked during the Medieval Climate Anomaly and declined during the last millennium, as climate became cooler and wetter and Native American burning declined. This trend was accentuated by 20th century fire suppression policies, which led to a minimum in burned area relative to the last 1400 years. In light of projected anthropogenic greenhouse gas emissions and predicted climate changes in California, our data indicate that although active management can mitigate local fire activity, broader regional burning may become more spatially extensive than has been observed in the last century.

Introduction

California's Camp Fire in 2018 was the state's most destructive wildfire on record, claiming 85 lives and damaging over 18 000 structures (CALFIRE 2019). These catastrophic events highlight a recent trend toward increasingly severe fire seasons and growing public concern, debate, and interest in fire ecology and policy in the Western United States (Hutto *et al* 2016, Schoennagel *et al* 2017). Fire-climate relationships lie at the heart of this concern as future climate change driven by anthropogenic greenhouse gas forcing is expected to bring warmer, drier conditions that promote more severe and destructive fire activity to this region (Westerling and Swetnam 2003, Van Wagtendonk and Lutz 2007, Barbero *et al* 2015). However, human management, and

in particular fire suppression, is also thought to have increased fire intensities at the expense of fire frequencies, due to the accumulation of fuel on previously managed landscapes. Understanding the 'top-down' controls of climate relative to the 'on-the-ground' influences of human activities on fire is crucial to developing effective management policies in an increasingly unstable earth system (Hunter 1996, Delcourt and Delcourt 1997, Keeley 2002). The influence and scales on which humans affect fire remain topics of debate, especially in North American ecosystems (Swetnam *et al* 2016, Syphard *et al* 2017, Roos *et al* 2018) where the impact of Native American peoples on fire regimes prior to the arrival of EuroAmerican colonists has important implications for our understanding of the long-term controls of fire (Caprio and Graber 2000).



In California, Native American fire use is undisputed, but its extent and influence on ecosystems is poorly understood (Parker 2002). Paleoecological records offer opportunities to investigate long-term fire-climate relationships, but discerning the role of humans in these fire histories is difficult as human burning cannot be easily differentiated from natural fire in paleoecological records (Whitlock *et al* 2010). We present a new analytical approach based on coeval measurements of two different sizes of charcoal to reconstruct and disentangle local and regional fire histories (Vachula *et al* 2018). Our approach represents a refinement of charcoal-based paleofire techniques and principles (Whitlock and Larsen 2002, Higuera *et al* 2007, Peters and Higuera 2007, Whitlock *et al* 2010, 2011). We focus on the central Sierra Nevada, an area known today for Yosemite National Park (figure 1). The region has rich cultural value among both the Miwok Native Americans that inhabited these mountains for millennia (Parker 2002), and the EuroAmericans who displaced these communities (Spence 1996). To characterize the relative influences of climate and human management on fire in the Sierra Nevada, we constructed a local and regional fire record using a sediment core from Swamp Lake, and compared our record with previously published records of regional temperature (Trouet *et al* 2013), moisture balance (Cook *et al* 1999, 2004), and human population density (Hull 2009).


Materials and methods

Sediment core and study site

A Bolivia core was taken from Swamp Lake (37.950 063°N, 119.829 074°, 1550 m a.s.l) in 2007 (Roach 2010, Denis *et al* 2012). Swamp Lake is located in the central Sierra Nevada Mountains and Yosemite National Park, California, USA. The lake has annually-laminated sediments and is relatively small (surface area = 0.08 km²) with seasonal inlet and outlet streams (Roach 2010, Street *et al* 2012). The core was split and imaged at LacCore with a Geotek Multi-Sensor Core Logger. Following Vachula *et al* (2018), minima of red and green color intensities from the Geotek high resolution images (1 pixel \approx 50 μ m) were counted manually to delineate annual laminations (Trauth 2007, Roach 2010). Additionally, dated macrovisual stratigraphic markers (a charcoal lamination, a slump sequence, and two tephtras) confirmed the varve-based age-depth chronology (table S1 is available online at stacks.iop.org/ERL/14/104011/mmedia).

Charcoal analysis

We followed standard procedures to isolate and identify charcoal particles (Whitlock and Larsen 2002, Chipman *et al* 2015). We volumetrically sampled (0.5–1.0 cm³) wet sediment from contiguous intervals



(1 cm) of the core and subjected these samples to a light chemical treatment (sodium hexametaphosphate and bleach; 24 h). Though our age-depth model resolution exceeded that of our sampling scheme, annual sampling was precluded by sediment availability and sampling accuracy. Next, we washed the samples over three nested sieves (63, 150, and 250 μm); with deionized water to isolate size fractions. Using a dissecting microscope (4–40 \times magnification), we counted charcoal particles in each fraction. Charcoal accumulation rates (CHARs) for each size fraction were calculated using the volumetric concentration and age-depth model (1):

$$\text{CHAR} = \frac{\# \text{ of Particles}}{\text{Volume}} * \frac{(\text{Top Depth} - \text{Bottom Depth})}{(\text{Top Age} - \text{Bottom Age})}.$$

In this letter, we consider the CHARs of two size fractions (63–150 and $>250 \mu\text{m}$, herein referred to as CHAR_{63-150} , and CHAR_{250} , respectively) to match and benefit from the source area determinations made by Vachula *et al* (2018). Vachula *et al* (2018) previously compared CHARs of different size fractions in Swamp Lake with historical fire data and identified the source area of CHAR_{250} and CHAR_{63-150} to be 25 and 150 km, respectively (Vachula *et al* 2018), thus providing a record of local ($>250 \mu\text{m}$, within 25 km of Swamp Lake) and regional (63–150 μm , within 150 km) area burned. Our new analytical approach allows us to differentiate fire histories of two distinct spatial scales. We compare these CHARs by calculating the linear correlation strength (Pearson's product moment correlation). To enable assessment of centennial-scale trends, raw CHARs were interpolated to the median sampling resolution of intervals (12 years) and smoothed (100 year lowess).


Results

The Swamp Lake record spanned 1431 years, as determined by counting annual laminations in high-resolution core images (figure S1). Our varve-based chronology agreed well with dated stratigraphic markers including a slump deposit, a visible lamination corresponding to a known fire event, and two tephra deposits (see supplementary text and table S1). To construct a fire history, we quantified the CHARs of two size fractions of sedimentary charcoal (63–150 and $>250 \mu\text{m}$).

Swamp Lake CHARs are inversely related to particle size; CHAR_{63-150} values (mean = $16.99 \text{ cm}^{-2} \text{ yr}^{-1}$; range = 0 to $47.44 \text{ cm}^{-2} \text{ yr}^{-1}$) were much greater than CHAR_{250} (mean = $0.54 \text{ cm}^{-2} \text{ yr}^{-1}$; range = 0 to $4.82 \text{ cm}^{-2} \text{ yr}^{-1}$; figure 2). Variations in CHARs in these three size fractions were generally positively correlated with one another throughout the record, though the strength of these correlations varied through time and at times the size fractions were inversely correlated. Correlation coefficients (200 year window) between CHAR_{63-150} and CHAR_{250} were greatest from ca. 850 to 1350 CE and ca. 1600 to 1900, whereas CHARs were less well correlated before ca. 850 CE and between ca. 1350 and 1600 CE, indicating varying correlations between local and regional fire history. This suggests temporal variations in the factors controlling local and regional fire.

Discussion


Both local and regional fire history is strongly influenced by climate, with warm and dry climates promoting burning. Area burned within 25 and 150 km was greater from 600 to 1500 CE, coincident with generally



warmer (Trouet *et al* 2013) and drier (Cook *et al* 1999, 2004) climates (figure 3). This finding is in line with previous work demonstrating that warm and dry climates promote burning in the Sierra Nevada (Brunelle and Anderson 2003), potentially via fuel desiccation and subsequently promoted fire ignition and spread. This conclusion disagrees with data suggesting wetter conditions promote fuel accumulation and widespread fires (Norman and Taylor 2003, Beaty and Taylor 2008, Vaillant and Stephens 2009). At centennial and shorter time-scales, climate-fire relationships are more convoluted, particularly at local spatial scales. Comparison of charcoal size classes indicates that although a drier climate from 900 to 1100 and 1200 to 1300 CE promoted both local and regional burning, the dry period from 1350 to 1550 only increased burning at regional scales (figure 3). Similarly, warm temperatures from 1000 to 1200 CE correspond with increased local and regional burning whereas warm temperatures from 700 to 900 and 1650 to 1750 CE appear to have promoted only regional scale burning (figure 3).

Whereas climate influenced both local and regional fire history, human land management and fire use

was a particularly significant control of local (<25 km) fire history in the central Sierra Nevada during intervals of the last millennium. The Sierra Miwok Native Americans inhabited this area and used fire to promote acorn crops from oak trees (Conrotto 1973, Anderson and Moratto 1996, Gassaway 2009), which served as their primary subsistence resource (McCarthy 1993, Stevens *et al* 2017). Reconstructions of Miwok population size based on debitage abundance in the Yosemite region also suggest that Native American populations varied considerably in size during the last 1400 years (Hull 2009). These population variations correspond with variations in burning at local scales (within 25 km) at times when the locally derived charcoal size fraction (CHAR₂₅₀) deviates from regionally sourced charcoal (CHAR_{63–150}). For example, periods of increased local burning corresponded with population maxima from 600 to 900, 1450 to 1700, and 1800 to 1900 CE (figure 3), with particularly distinct state-changes in the Miwok population and fire from 700 to 850, 1475–1525, and 1700 to 1850 CE. These relationships suggest that humans influenced local fire history (<25 km) and caused disconnects between climate and fire, and in the spatial homogeneity of fire history.



The spatial self-similarity, coupling, and decoupling of local and regional fire history evident in the Swamp Lake record are intriguing phenomena. Conceptual models have postulated that the controls of fire are self-similar across a range of spatial and temporal scales (Parisien and Moritz 2009, Whitlock *et al* 2010). Though some research has sought to characterize the cross-scale dynamics of fire using modern observational data (Díaz-Delgado *et al* 2004, Falk *et al* 2007, Slocum *et al* 2010), relatively little has been done to assess these dynamics on the longer timescales relevant to climate change (Scholl and Taylor 2010, Whitlock *et al* 2010). Our record indicates that (1) climate changes influenced burning at all spatial scales, (2) Native American influences appear to have been limited to local scales, but (3) high Miwok populations resulted in fire even during periods of climate conditions unfavorable to fires (figure 3). Thus, our record indicates that human alteration of natural fire regimes causes disconnects in the areal self-similarity of fire. Climate, on the other hand, acts as a ‘top-down’ control on regional fire, which, in the absence of human intervention, cascades down to local scales of fire dynamics. In sum, these cross-scale relationships

suggest human management acts as a bottom-up control of fire that can interact with, override, or be overridden by top-down climatic influences.

Both CHAR₂₅₀ and CHAR_{63–150} exhibit steady declines over the last 1400 years, in tandem with the development of cool and wet conditions not conducive to burning. These long-term decreasing trends in local and regional fire culminate the 20th century, which experienced the lowest levels of fire activity in the last 1400 years (figure 3). Federal policies suppressed burning in the Sierra Nevada from ca. 1900 to 1970 CE (Taylor and Scholl 2012). This period of suppression is clearly evident in our charcoal record (figure 4). The suppression period experienced warmer and drier climate conditions than the preceding century, yet burning was muted at both regional and local scales (figure 4). This disconnect is thought to have resulted in a subsequent fuel surplus, which has promoted recent burning following the cessation of suppression policies (Caprio and Graber 2000, Westerling *et al* 2003, Miller *et al* 2009, Steel *et al* 2015). However, our data demonstrate recent burning is only remarkable in how limited it has been in the context of the last 1400 years (figure 3). This finding

implies that fires have the potential and the precedent to be much more spatially extensive than we have observed in the last century.

Our data suggest that though human management can influence local fire, a warmer and drier climate controls large-scale area burned. The fire-climate relationships elucidated from our dataset, in combination with projections for warmer and drier climate conditions in California (Hayhoe *et al* 2004), suggest that future burning in the Sierra Nevada will be more spatially extensive than has been observed in the last century. The legacy of fire suppression, which is clearly evident in our data (figure 3), could further exacerbate fires of the future (Donovan and Brown 2007).

Acknowledgments

This work was supported by a graduate student fellowship and research grant to RSV by the Institute at Brown for Environment and Society. This work was also supported by the National Park Service Shared Beringia Heritage program, Department of the Interior-P18AC00556. We thank A Cheung for helpful discussions and K Brady and LacCore for sampling assistance.

Data availability statement

The charcoal data that support the findings of this study are included within the article. The PDSI (Cook *et al* 1999, 2004) and paleotemperature (Trouet *et al* 2013) data are available from the World Data Center for Paleoclimatology: <https://ncdc.noaa.gov/paleo>. The Yosemite National Park visitor data are available from the National Park Service data store: <https://irma.nps.gov>. The Yosemite annual temperature data are available from the NOAA National Climatic Data Centers: <https://ncdc.noaa.gov>.

ORCID iDs

Richard S Vachula  <https://orcid.org/0000-0001-5559-6540>

References

- Anderson M K and Moratto M J 1996 Native American land-use practices and ecological impacts *Sierra Nevada Ecosystem Project: Final Report to Congress* vol 2 University of California, Center for Water and Wildland Resources Davis pp 187–206
- Barbero R, Abatzoglou J T, Larkin N K, Kolden C A and Stocks B 2015 Climate change presents increased potential for very large fires in the contiguous United States *Int. J. Wildlife Fire* **24** 892
- Beaty R M and Taylor A H 2008 Fire history and the structure and dynamics of a mixed conifer forest landscape in the northern Sierra Nevada, Lake Tahoe Basin, California, USA *For. Ecol. Manage.* **255** 707–19
- Brunelle A and Anderson R S 2003 Sedimentary charcoal as an indicator of late-Holocene drought in the Sierra Nevada, California, and its relevance to the future *Holocene* **13** 21–8
- CALFIRE 2019 *Top 20 Most Destructive California Wildfires* (http://fire.ca.gov/communications/downloads/fact_sheets/Top20_Destruction.pdf)
- Caprio A C and Graber D M 2000 Returning fire to the mountains: can we successfully restore the ecological role of pre-Euroamerican fire regimes to the Sierra Nevada? *Wilderness Sci. Time Chang. Conf.* **5** 233–41
- Chipman M L, Hudspeth V, Higuera P E, Duffy P A, Kelly R, Oswald W W and Hu F S 2015 Spatiotemporal patterns of tundra fires: late-Quaternary charcoal records from Alaska *Biogeosciences* **12** 4017–27
- Conrotto E L 1973 *Miwok Means People: the Life and Fate of the Native Inhabitants of the California Gold Rush Country* (Fresno, CA: Valley Publishers)
- Cook E R, Meko D M, Stahle D W and Cleaveland M K 1999 Drought reconstructions for the continental United States *J. Clim.* **12** 1145–62
- Cook E R, Woodhouse C A, Eakin C M, Meko D M and Stahle D W 2004 Long-term aridity changes in the western United States *Science* **306** 1015–8
- Delcourt H R and Delcourt P A 1997 Pre-Columbian native American use of fire on southern Appalachian landscapes *Conserv. Biol.* **11** 1010–4
- Denis E H, Toney J L, Tarozo R, Scott Anderson R, Roach L D and Huang Y 2012 Polycyclic aromatic hydrocarbons (PAHs) in lake sediments record historic fire events: validation using HPLC-fluorescence detection *Org. Geochem.* **45** 7–17
- Díaz-Delgado R, Lloret F and Pons X 2004 Spatial patterns of fire occurrence in Catalonia, NE, Spain *Landsc. Ecol.* **19** 731–45
- Donovan G H and Brown T C 2007 Be careful what you wish for: the legacy of Smokey Bear *Front. Ecol. Environ.* **5** 73–9
- Falk D A, Miller C, McKenzie D and Black A E 2007 Cross-scale analysis of fire regimes *Ecosystems* **10** 809–23
- Gassaway L 2009 Native American fire patterns in Yosemite Valley: archaeology, dendrochronology, subsistence, and culture change in the Sierra Nevada *Soc. Calif. Archaeol. Proc.* **22** 1–19
- Hayhoe K *et al* 2004 Emissions pathways, climate change, and impacts on California *Proc. Natl Acad. Sci. USA* **101** 12422–7
- Higuera P E, Peters M E, Brubaker L B and Gavin D G 2007 Understanding the origin and analysis of sediment-charcoal records with a simulation model *Quat. Sci. Rev.* **26** 1790–809
- Higuera P E, Whitlock C and Gage J A 2011 Linking tree-ring and sediment-charcoal records to reconstruct fire occurrence and area burned in subalpine forests of yellowstone National Park, USA *Holocene* **21** 327–41
- Hull K L 2009 *Pestilence and Persistence: Yosemite Indian Demography and Culture in Colonial California* (Berkeley: University of California Press)
- Hunter M 1996 Benchmarks for managing ecosystems: are human activities natural? *Conserv. Biol.* **10** 695–7
- Hutto R L, Keane R E, Sherriff R L, Rota C T, Eby L A and Saab V A 2016 Toward a more ecologically informed view of severe forest fires *Ecosphere* **7** e01255
- Keeley J E 2002 Fire management of California shrubland landscapes *Environ. Manage.* **29** 395–408
- McCarthy H 1993 Managing oaks and the acorn crop *Before Wilderness Environ. Manag. by Nativ. Californians* pp 213–28
- Miller J D, Safford H D, Crimmins M and Thode A E 2009 Quantitative evidence for increasing forest fire severity in the sierra nevada and southern cascade mountains, California and Nevada, USA *Ecosystems* **12** 16–32
- National Park Service 2018 Annual Park Recreation Visitation 1904–2018 ([https://irma.nps.gov/Stats/SSRSReports/ParkSpecificReports/AnnualParkRecreationVisitation\(1904—LastCalendarYear\)?Park=YOSE](https://irma.nps.gov/Stats/SSRSReports/ParkSpecificReports/AnnualParkRecreationVisitation(1904—LastCalendarYear)?Park=YOSE))
- NOAA 2019 NNDC CLIMATE DATA ONLINE *Satell. Inf. Serv*
- Norman S P and Taylor A H 2003 Tropical and north Pacific teleconnections influence fire regimes in pine-dominated forests of north-eastern California, USA *J. Biogeogr.* **30** 1081–92
- Parisien M-A and Moritz M A 2009 Environmental controls on the distribution of wildfire at multiple spatial scales *Ecol. Monogr.* **79** 127–54

- Parker A J 2002 Fire in Sierra Nevada forests: evaluating the ecological impact of burning by Native Americans *Fire, Nativ. Peoples, Nat. Landsc.* pp 233–67
- Peters M E and Higuera P E 2007 Quantifying the source area of macroscopic charcoal with a particle dispersal model *Quat. Res.* **67** 304–10
- Roach L D 2010 *Climate Change in the Pacific North America Region Over the Past Millennium: Development and Application of Novel Geochemical Tracers* (San Diego, CA: University of California)
- Roos C I, Zedeño M N, Hollenback K L and Erlick M M H 2018 Indigenous impacts on North American Great Plains fire regimes of the past millennium *Proc. Natl Acad. Sci.* **115** 8143–8
- Schoennagel T, Balch J K, Brenkert-Smith H, Dennison P E, Harvey B J, Krawchuk M A, Mietkiewicz N, Morgan P, Moritz M A and Rasker R 2017 Adapt to more wildfire in western North American forests as climate changes *Proc. Natl Acad. Sci.* **114** 4582–90
- Scholl A E and Taylor A H 2010 Fire regimes, forest change, and self-organization in an old-growth mixed-conifer forest, Yosemite National Park, USA *Ecol. Appl.* **20** 362–80
- Slocum M G, Beckage B, Platt W J, Orzell S L and Taylor W 2010 Effect of climate on wildfire size: a cross-scale analysis *Ecosystems* **13** 828–40
- Spence M 1996 Dispossessing the Wilderness: Yosemite Indians and the National Park Ideal, 1864–1930 *Pac. Hist. Rev.* **65** 27–59
- Steel Z L, Safford H D and Viers J H 2015 The fire frequency-severity relationship and the legacy of fire suppression in California forests *Ecosphere* **6** 8
- Stevens N E, Whitaker A R and Rosenthal J S 2017 Bedrock mortars as indicators of territorial behavior in the Sierra Nevada *Quat. Int.* **518** 57–68
- Street J H, Anderson R S and Paytan A 2012 An organic geochemical record of Sierra Nevada climate since the LGM from Swamp Lake, Yosemite *Quat. Sci. Rev.* **40** 89–106
- Swetnam T W, Farella J, Roos C I, Liebmann M J, Falk D A and Allen C D 2016 Multiscale perspectives of fire, climate and humans in western North America and the Jemez Mountains, USA *Phil. Trans. R. Soc. B* **371** 20150168
- Syphard A D, Keeley J E, Pfaff A H and Ferschweiler K 2017 Human presence diminishes the importance of climate in driving fire activity across the United States *Proc. Natl Acad. Sci.* **114** 13750–5
- Taylor A H and Scholl A E 2012 Climatic and human influences on fire regimes in mixed conifer forests in Yosemite National Park, USA *For. Ecol. Manage.* **267** 144–56
- Trauth M H 2007 *MATLAB Recipes for Earth Sciences* ed M H Trauth (Berlin: Springer)
- Trouet V, Diaz H F, Wahl E R, Viau A E, Graham R, Graham N and Cook E R 2013 A 1500-year reconstruction of annual mean temperature for temperate North America on decadal-to-multidecadal time scales *Environ. Res. Lett.* **8** 24008
- Vachula R S, Russell J M, Huang Y and Richter N 2018 Assessing the spatial fidelity of sedimentary charcoal size fractions as fire history proxies with a high-resolution sediment record and historical data *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **508** 166–75
- Vaillant N M and Stephens S L 2009 Fire history of a lower elevation Jeffrey pine-mixed conifer forest in the eastern Sierra Nevada, California *USA Fire Ecol.* **5** 4–19
- Van Wagtenonk J W and Lutz J A 2007 Fire regime attributes of wildland fires in Yosemite National Park *USA Fire Ecol.* **3** 34–52
- Westerling A L, Hidalgo H G, Cayan D R and Swetnam T W 2003 Warming and earlier spring increase western U.S. Forest Wildfire Activity *Science* **400** 1054
- Westerling A L and Swetnam T W 2003 Interannual to decadal drought and wildfire in the western United States *EOS Trans. Am. Geophys. Union* **84** 545–55
- Whitlock C, Higuera P E, McWethy D B and Briles C E 2010 Paleocological perspectives on fire ecology: revisiting the fire-regime concept ~!2009-09-02~!2009-11-09~!2010-03-05~! *Open Ecol. J.* **3** 6–23
- Whitlock C and Larsen C 2002 Charcoal as a fire proxy *Tracking Environmental Change Using Lake Sediments* (Dordrecht: Kluwer) pp 75–97